

Twenty-three polymorphic microsatellite markers for the Caribbean endemic *Zenaida* dove, *Zenaida aurita*, and its conservation in related *Zenaida* species

Karine Monceau · Maria Gaillard · Estelle Harrang ·
Diego Santiago-Alarcon · Patricia G. Parker ·
Frank Cezilly · Rémi A. Wattier

Received: 10 December 2008 / Accepted: 16 December 2008 / Published online: 4 January 2009
© Springer Science+Business Media B.V. 2009

Abstract Twenty-three polymorphic microsatellite loci, six dinucleotidic loci and 17 tetranucleotidic loci, were developed for the *Zenaida* dove (*Zenaida aurita*), a bird species endemic to the Caribbean Islands. From a set of 30 individuals captured at one single location in Barbados, we obtained 20 loci that did not deviate from Hardy–Weinberg Equilibrium. Number of alleles per locus ranged between 2 and 11 (average 7.05) and the expected heterozygosity per locus, H_e ranged between 0.321 and 0.881 (average 0.712). This gives an exclusionary power for parental analysis of 0.9999 and 1.0000, knowing the genotype of one social parent, or both, respectively. Such results indicate that these 20 loci will be useful for both studying population genetics and mate choice patterns in *Z. aurita*. All 20 loci amplified in four other *Zenaida* species, the Galápagos dove, *Z. galapagoensis*, the eared dove, *Z. auriculata*, the mourning dove, *Z. macroura*, the Pacific dove, *Z. meloda*, with 30–96% being polymorphic.

Keywords Endemic species · Extra-pair paternity · Gene flow · Mate choice · Microsatellite · *Zenaida* dove

K. Monceau · M. Gaillard · E. Harrang · F. Cezilly ·
R. A. Wattier (✉)
Equipe Ecologie Evolutive, UMR CNRS 5561 Biogéosciences,
Université de Bourgogne, 6 Boulevard Gabriel, 21000 Dijon,
France
e-mail: Remi.wattier@u-bourgogne.fr

D. Santiago-Alarcon · Patricia G. Parker
Department of Biology, University of Missouri-St. Louis,
One University Boulevard, Saint Louis, MO 63121, USA

Patricia G. Parker
Wild Care Institute, Saint Louis Zoo, 1 Government Drive,
St. Louis, MO 63110, USA

The *Zenaida* dove (*Zenaida aurita*) is a columbid species endemic to the Caribbean Islands (Wiley 1991). This species is of patrimonial interest, and is legally hunted in Puerto Rico, Guadeloupe and Martinique. It suffers in some islands from competition with the invasive Eurasian collared dove (*Streptopelia decaocto*), habitat destruction, and illegal hunting (Barré et al. 1997; Rivera-Milan 1997), very little is known, of movements of individuals between islands. In addition, knowledge of the mating system of the *Zenaida* dove remains limited. Although the species is territorial and socially monogamous (Wiley 1991), the occurrence of extra-pair paternity has not been investigated so far. However, data on mate fidelity and extra-pair paternity in tropical species that maintain year-round pair bonds is particularly valuable, yet rare (Stutchbury and Morton 2001).

Molecular markers are needed to both assess the conservation genetics of this species (i.e. allelic diversity related to island size, level of gene flow, and differentiation) and study mate choice and mating patterns (i.e. female choice related to male heterozygosity and genetic similarity, and rates of extra-pair paternity).

Four *Zenaida aurita* partial genomic DNA libraries enriched for microsatellite sequences containing CA, GA, CATC and TAGA repeat motifs were constructed following the methods described in Jones et al. (2002). High molecular weight DNA was extracted from blood using chloroform/phenol/isopropanol extraction (Sambrook et al. 1989) from 10 individuals from one location in Barbados. DNA was partially digested with a cocktail of seven blunt-end restriction enzymes (*Rsa*I, *Hae*III, *Bsr* BI, *Pvu*II, *Stu*I, *Sca*I and *Eco* RV). Fragments 350–700 bp long were adapted with 20 bp oligonucleotides which contained a *Hind*III site at the 5' end, and subjected to magnetic bead capture using 5'-biotinylated microsatellite oligonucleotides as capture

Table 1 Characterization of 23 polymorphic microsatellite loci for Zenaida dove (*Zenaida aurita*)

Locus	Repeat motif	Primer sequence (5'-3')	Genbank Accession #	Size Range (bp)	N	N _A	H _o	H _e	PE-1	PE-2
ZaA4	(GT) ₁₉	F: AGCAGGCTCATTAACACATTAG R: ACCTGTAGTTCAGGGCGTATAG	GF091744	243–273	29	6	0.897	0.796	0.396	0.575
ZaA5	(GT) ₂₁	F: GGCACACAAATACTTTAAGAGG R: GACCTTTGCTCACTCACTATTC	GF091766	251–285	29	11	0.724	0.848	0.501	0.670
ZaA7	(AC) ₂₄	F: TGGCACTGAAGAACAAGC R: CTGGGACTCGGGATTAC	GF091745	101–147	26	9	0.769	0.759	0.367	0.551
ZaA112	(AC) ₁₉	F: CTCCTGGTTTCAITTCAGAC R: ATGGTGCATTTGTGTAGG	GF091746	194–204	29	6	0.862	0.736	0.310	0.485
ZaA113	(AC) ₂₁	F: AGCTTTTGATAAAGCCTGAAC R: CTTGCTGTATGCCACAATG	GF091747	210–248	29	9	0.828	0.872	0.549	0.712
ZaA127	(CA) ₁₇	F: GCATTTGCTCCTCCAGTAGATT R: AAGCCTGACAGTATTCAGAAG	GF091748	245–247	30	2	0.333	0.427	0.088	0.166
ZaC1	(TCCA) ₉	F: TCCTGAGCTTCTGGGACAC R: GGGACAAGGAGCATCCAC	GF091749	174–194	29	4	0.586	0.607	0.187	0.344
ZaC3	(TCCA) ₁₀	F: TTCTTCTGCCAGCACTGTG R: TTTTCCCTACCTGCCCTTTTC	GF091750	252–272	28	4	0.429	0.459	0.103	0.238
ZaC11	(TGGA) ₄ (GGGA)(TGGG) ₄	F: GCTTTCATCTGTGTGAGGAC R: AGGACATCCATAGCCTGATAC	GF091751	123–151	29	5	0.828	0.791	0.386	0.565
ZaC12	(TCCA) ₁₂	F: AGCTCCGAACCTCTAACTAC R: GATAGGCTCTCTGCTGATG	GF091752	279–323	30	9	0.700	0.788	0.400	0.579
ZaC117	(GGAT) ₁₁	F: AAGGCAAAAAGGAAAGGAAAG R: TCCGTGATTTATGCTGTAGC	GF091753	128–156	27	7	0.852	0.665	0.246	0.412
ZaD1	(TAGA) ₁₁	F: TAGCGTGGACAGACTTGTG R: GGGTGAITTTGATGTTCCCTTC	GF091754	121–165	29	8	0.759	0.846	0.492	0.664
ZaD5	(TATC) ₁₃	F: ACCACCCCTCAITTTGTAAGAT R: ACACACACAGCAGTAAGTTGAC	GF091755	162–282	29	7	0.207*	0.797	–	–
ZaD6	(TAGA) ₈	F: TCTTGACTTCACTCCTGACAC R: CCTACTCCACATCTTCTGTT	GF091756	241–269	30	6	0.500	0.548	0.163	0.330
ZaD7	(TCTA) ₄ (TCCTAA)(TCTA) ₉	F: CCTTTTGAAGGTGTTCAATC R: GCATCCAAAATGAGTATGC	GF091757	304–332	30	8	0.833	0.833	0.477	0.651
ZaD10	(TTCTA) ₈ (TTCITG) ₄ N ₂ (TCCTA) ₁₆	F: CGAGCCTGTCAGTGTTC R: TCATGGGTTACTGTTACAACCTC	GF091758	368–416	30	11	0.400*	0.894	–	–
ZaD11	(TCTA) ₁₅	F: TGAACCATTTGCTCCAGAG R: TGCCATAACTTTGAGAACTGC	GF091759	230–266	28	7	0.893	0.803	0.418	0.596
ZaD104	(GATA) ₁₀	F: ACCGAAAATCCTAACCTGAAC R: ATGCCAGTCTGTGAAAACAAG	GF091760	245–273	28	7	0.929	0.841	0.477	0.651

Table 1 continued

Locus	Repeat motif	Primer sequence (5'–3')	Genbank Accession #	Size Range (bp)	N	N _A	H _o	H _e	PE-1	PE-2
ZaD105	(GATA) ₁₁	F: GCAGGAATCAGCTTTCAGAGTAC R: TTGAAGATGATACAGCAGAGGTG	GF091761	264–284	30	6	0.733	0.647	0.226	0.388
ZaD108	(TCTA) ₁₀	F: CCTTCCCTTCAATGACTTACC R: GCATCTGTGTCATCCAG	GF091762	141–173	30	9	0.833	0.881	0.571	0.729
ZaD116	(CTAT) ₁₃	F: CAGGGGAAAAAAAATCT R: TTCTGAGTGAAGATGGGTAAT	GF091763	123–143	28	6	0.321*	0.761	–	–
ZaD119	(GATA) ₁₂	F: CATTAACGCAATGTTTCTGAG R: TGCCCAAAGGTTTGTATTTAC	GF091764	162–186	29	7	0.897	0.758	0.353	0.531
ZaD121	(GATA) ₁₃	F: CTGAGGTTCCAACAACACTTG R: TTAGAGGACTGTCGTTCTGTG	GF091765	203–247	29	10	0.897	0.909	0.641	0.783
Overall									0.999	1.000

N = Number of individual typed, all coming from the same location in Barbados. N_A = Number of alleles per locus. H_o and H_e = Observed and expected heterozygotes, respectively. PE-1 and PE-2 are exclusion probabilities of knowing the genotype of one or both parents, respectively

* Significant deviation from Hardy–Weinberg equilibrium after Bonferroni correction for multiple tests (P < 0.00212)

molecules. Fragments enriched for microsatellites were subsequently PCR amplified using primer complementary to the adaptor, digested with *HindIII* to remove the adaptor sequences, and ligated into the *HindIII* site of pUC19. The plasmids were then electroporated into *Escherichia coli* DH5 α . Recombinant clones, identified by white–blue screening, were chosen at random for sequencing on an ABI373 automated sequencer, using ABI Prism Taq dye terminator cycle sequencing methodology. A total of 81 out of 94 positive clones sequenced contained microsatellites (86.17%). Primers were designed for 50 microsatellite-bearing clones using Designer PCR, versus 1.03 (Research Genetics, Inc.). Enriched library construction, sequencing of positive clones and primer design were performed by Genetic Identification Services (www.genetic-id-services.com). Primers pairs were synthesized (GENCUST, France) for 33 potential loci, each forward primer being 5' tailed with a M13 sequence (5'-CACGACGTTGTAAAACGAC-3'). PCRs were carried out in 10 μ l volume including 5 ng DNA template, 200 μ M each dNTP, 200 nM each primer (Table 1), 0.025 μ M of 5' labelled M13 primer with 700 and 800 fluorochromes (LICOR), 1 \times reaction buffer and 0.15 U HotMaster DNA polymerase (5-PRIME). A T3 thermocycler (Biometra) was used beginning with an initial denaturation at 94°C for 1.5 min, followed by 35 cycles consisting of 20 s at 94°C, 30 s at 46°C and 30 s at 65°C, and a final extension step at 65°C for 3 min. All 33 loci successfully amplified as tested on 6 cm long 2% agarose gels (TBE 1 \times) for 30 min at 100 V in a RunOne electrophoresis chambers (EmbiTec). Product size variations were visualized using a LICOR 4000L automated sequencer, 6.5% acrylamide, 41 cm long gels. Image analysis and allele scoring were carried out by eye by two independent investigators and then compared. Ten primer pairs either failed to reveal any polymorphism or produced unreadable patterns on a set of eight individuals from one location in Barbados. Polymorphism of the 23 remaining loci was assessed on an extra set of up to 22 Zenaida doves from the same location. Estimates of observed (H_o) and expected (H_e) heterozygosity, deviations from Hardy–Weinberg equilibrium (HWE), and genotypic linkage disequilibrium (LD) were estimated using the software FSTAT version 2.9.3.2 (Goudet 1995). Exclusion probabilities knowing the genotype of one or both social parents, PE-1 and PE-2, respectively, were estimated for loci not deviating significantly from HWE using the software CERVUS 2.0 (Marshall et al. 1998).

No linkage disequilibrium was detected after Bonferroni correction for multiple tests. Three loci ZaD5, ZaD10 and ZaD116 showed a heterozygote deficiency, significantly deviating (after Bonferroni correction) from Hardy–Weinberg Equilibrium (Table 1) probably due to null alleles associated with absence of conservation of flanking

Table 2 Trans-specific test of amplification and polymorphism of 23 microsatellite loci initially developed for *Zenaida* dove (*Zenaida aurita*) for four other *Zenaida* species and one *Colombina* species

Locus	<i>Z. galapagoensis</i> (n = 3)	<i>Z. auriculata</i> (n = 3)	<i>Z. macroura</i> (n = 1)	<i>Z. meloda</i> (n = 3)	<i>Colombina passerina</i> (n = 2)
ZaA4	1	3	1	2	1
ZaA5	2	2	1	1	–
ZaA7	1	2	2	1	–
ZaA112	1	2	1	1	1
ZaA113	4	3	1	1	–
ZaA127	2	3	2	2	1
ZaC1	2	2	1	4	2
ZaC3	4	3	1	2	1
ZaC11	2	4	1	1	3
ZaC12	4	4	1	–	1
ZaC117	4	3	2	1	–
ZaD1	2	3	2	4	–
ZaD5	1	2	2	2	–
ZaD6	4	3	2	2	1
ZaD7	4	3	1	3	1
ZaD10	2	2	1	–	1
ZaD11	6	4	1	4	–
ZaD104	1	4	1	1	1
ZaD105	3	2	1	1	1
ZaD108	1	1	2	1	1
ZaD116	2	3	1	–	–
ZaD119	2	2	1	2	1
ZaD121	5	4	1	1	1
AS (%)	100	100	100	87	65
P (%)	74	96	30	50	13
Na	3.18	2.86	2.00	2.70	2.50

N = Number of individuals tested. – Denotes an absence of amplification.
AS = Amplification success, percentage of amplifying loci.
P = Percentage of polymorphic loci. Na = Mean number of alleles for polymorphic loci

sequences (Wattier et al. 1998). Therefore, we recommend not using these loci. The remaining 20 loci consist of 2–11 alleles (average 7.05), an expected heterozygosity (H_e) ranging from 0.427 to 0.909 (average 0.712) and an exclusionary power of 0.9999 and 1.0000, knowing the genotype of one social parent, or both, respectively (Table 1). This exclusionary power remains extremely high if only the eight loci with the highest individual PE-1 are taken into account, combined PE-1 = 0.9972 and PE-2 = 0.9999. These results indicate that the 20 polymorphic loci not deviating from HWE will be useful for both studying population genetics and mate choice patterns in *Z. aurita*.

Amplification success and polymorphism were assessed for four other *Zenaida* species (Table 2): the Galápagos dove, *Z. galapagoensis* (three from Galápagos), the eared dove, *Z. auriculata* (two from Uruguay and one from Venezuela), the mourning dove, *Z. macroura* (one individual from the USA), the Pacific dove, *Z. meloda* (three from Peru) and one *Colombina* species, the common ground-dove, *Colombina passerina*. (one from Mexico and one from Venezuela). PCRs and genotyping conditions

were identical to those used for *Z. aurita*. The overall success of amplification (Table 2) for the four *Zenaida* species is extremely high (87–100%), being 100% for the three most related to *Z. aurita*, that is *Z. galapagoensis*, *Z. auriculata* and *Z. macroura* (Johnson and Clayton 2000). Many loci are polymorphic (30–96%) with 2.00–3.18 alleles per loci on average (Table 2). These results are very encouraging about the usefulness of the loci developed in *Z. aurita* in other *Zenaida* species, especially taking into account the limited sampling size used ($n = 3$ for three species and only one for *Z. macroura*). Until now, only five loci were available for one species, the Galápagos dove (Santiago-Alarcon et al. 2006). From the 65% of the loci amplified in *C. passerina*, 13% were polymorphic with 2.5 alleles on average. These results also show that the loci developed for *Z. aurita* are likely to be useful for other Columbidae.

Acknowledgements This work was supported by grants to F. C. and R. W. by “Institut Buffon” Université de Bourgogne, to K. M. from the French Ministère de l’Éducation Nationale (grant # 01121) and to D. S.-A. and P. G. P. from the Whitney R. Harris World Ecology Centre, Idea Wild, The Frank M. Chapman Memorial Fund of the

American Museum of Natural History, The Field Research for Conservation program of the Saint Louis Zoo (FRC 05-2 and FRC 08-2), the Organization for Tropical Studies, and by the Des Lee collaborative vision in zoological studies. Logistical support and permits were provided by the Galápagos National Park, the Darwin Research Foundation and the Nature Heritage Department of Barbados. Authors thank Kenneth Jones and Mark Todd from Genetic Identification Services for their help and patience in developing the microsatellite loci, and Jérôme Moreau for helping collecting biological material.

References

- Barré N, Feldmann P, Tayalay G et al (1997) Introduction et extension de la tourterelle turque *Streptopelia decaocto* dans les Petites-Antilles. *Alauda* 65:245–250
- Goudet J (1995) FSTAT (vers. 1.2): a computer program to calculate F-statistics. *J Hered* 86:485–486
- Johnson KP, Clayton DH (2000) A molecular phylogeny of the dove genus *Zenaida*: mitochondrial and nuclear sequences. *Condor* 102:864–870. doi:10.1650/0010-5422(2000)102[0864:AMPOTD]2.0.CO;2
- Jones KC, Levine KF, Banks JD (2002) Characterization of 11 polymorphic tetranucleotide microsatellites for forensic applications in California elk (*Cervus elapus canadensis*). *Mol Ecol Notes* 2:425–427. doi:10.1046/j.1471-8286.2002.00264.x
- Marshall TC, Slate J, Kruuk L et al (1998) Statistical confidence for likelihood-based paternity inference in natural populations. *Mol Ecol* 7:639–655. doi:10.1046/j.1365-294x.1998.00374.x
- Rivera-Milan FF (1997) Seasonal and annual changes in the population density of Zenaida doves in the xenophytic forest of Guanica, Puerto Rico. *J Field Ornithol* 68:259–272
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning: a laboratory manual*, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Santiago-Alarcon D, Tanksley SM, Parker PG (2006) Morphological variation and genetic structure of the Galápagos dove (*Zenaida galapagoensis*) populations: issues in conservation for the Galápagos bird fauna. *Wilson J Ornithol* 118:194–207. doi:10.1676/05-010.1
- Stutchbury BJM, Morton ES (2001) *Behavioural ecology of tropical birds*. Academic Press, New York
- Wattier R, Engel CR, Saumitou-Laprade P et al (1998) Short allele dominance as a source of heterozygote deficiency at microsatellite loci: experimental evidence at the dinucleotide locus Gv1CT in *Gracilaria gracilis* (Rhodophyta). *Mol Ecol* 7:1569–1573. doi:10.1046/j.1365-294x.1998.00477.x
- Wiley JW (1991) Ecology and behaviour of the Zenaida dove. *Ornitol Neotrop* 2:49–75